

Synthesis of coumarin or ferrocene labeled nucleosides via Staudinger ligation

Ivana Kosiova*, Andrea Janicova and Pavol Kois

Full Research Paper

Open Access

Address:
Comenius University, Faculty of Natural Sciences, Department of
Organic Chemistry, Mlynska dolina, Pavilon CH2, SK-84215
Bratislava, Slovak Republic

Email:
Ivana Kosiova* - kosiova@fns.uniba.sk; Andrea Janicova -
janicova@fns.uniba.sk; Pavol Kois - kois@fns.uniba.sk

* Corresponding author

Beilstein Journal of Organic Chemistry **2006**, 2, No. 23.
doi:10.1186/1860-5397-2-23

Received: 16 October 2006
Accepted: 30 November 2006
Published: 30 November 2006

© 2006 Kosiova et al; licensee Beilstein-Institut.
License and terms: see end of document.

Abstract

Background

Reaction of azides with triaryl phosphines under mild conditions gives iminophosphoranes which can react with almost any kind of electrophilic reagent, e.g. aldehydes/ketones to form imines or esters to form amides. This so-called Staudinger ligation has been employed in a wide range of applications as a general tool for bioconjugation including specific labeling of nucleic acids.

Results

A new approach for the preparation of labeled nucleosides via intermolecular Staudinger ligation is described. Reaction of azido-nucleosides with triphenylphosphine lead to iminophosphorane intermediates, which react subsequently with derivatives of coumarin or ferrocene to form coumarin or ferrocene labeled nucleosides. Fluorescent properties of coumarin labeled nucleosides are determined.

Conclusion

New coumarin and ferrocene labeled nucleosides were prepared *via* intermolecular Staudinger ligation. This reaction joins the fluorescent coumarin and biospecific nucleoside to the new molecule with promising fluorescent and electrochemical properties. The isolated yields of products depend on the structure of azidonucleoside and carboxylic acids. A detailed study of the kinetics of the Staudinger ligation with nucleoside substrates is in progress.

Background

Modified nucleosides are important tools for the study of key processes of cell metabolism, as well as being successful therapeutic agents. [1-3] For structural and functional studies of

nucleosides, and their oligomers in nucleic acids or protein complexes, efficient detection methods are necessary. Today, fluorescent detection is of paramount importance to biological

studies. [4-10] The sensitivity of fluorescence techniques has reached an extremely high level, similar to radioactive methods, and can even provide information on the dynamic structure of dye-bound biomolecules. Complicated and expensive optical detection would be gradually replaced by simpler and cheaper electrochemical detection based on the redox properties and electrical conductance of biomolecules. [11,12] Electrochemical techniques can be highly sensitive, rapid and available to production in miniaturised formats. Although different approaches for modification of nucleic acid components and DNA biosensor construction have been developed, many questions remain to be answered with respect to the complete understanding of optical and electrical properties of modified nucleic acids used in bioanalytical systems.

Universal starting compounds for preparation of modified nucleosides are azidonucleosides. In general, organic azides are valuable, energy-rich and flexible intermediates, which can react very differently under various reaction conditions. [13] They can react at N1 with electrophiles (carbon electrophiles, protons, boranes) and at N3 with nucleophiles, very frequently with phosphorous nucleophiles. Reaction of azides with triaryl phosphines under mild conditions gives iminophosphoranes without formation of any byproducts. [14] The intermediate which is formed almost quantitatively can be rapidly hydrolysed to the primary amine and triarylphosphine oxide. This Staudinger reduction is a frequently used method for the smooth reduction of azides to amines. Iminophosphoranes can react with almost any kind of electrophilic reagent, [15,16] e.g. aldehydes or ketones to form imines. Also less reactive carbonyl electrophiles, such as esters, can undergo reaction with iminophosphorane to form amides, especially if the electrophilic attack proceeds in an intramolecular fashion. [17-21] This so-called Staudinger ligation has been employed in a wide range of applications as a general tool for bioconjugation, [22,

23] including specific labeling of nucleic acids, [24] proteomic studies [25,26] and modification of cell surfaces. [17,18]

We applied the Staudinger ligation for nucleoside labeling procedures, using coumarin and ferrocene derivatives as labels. According to our knowledge, applications of this reaction in nucleoside and nucleotide chemistry are rare.


Results and discussion

We used intermolecular Staudinger ligation [27] for the preparation of fluorescent and electrochemically labeled nucleosides. As the label we used either coumarin-4-acetic acids **1a-c** or ferrocene derivative **17** (Figure 1), which were synthesised in our department as part of studies into new ligands. Coumarins as fluorescent probes or labels [28-30] have extensive and diverse applications, they exhibit extended spectral range, are photostable and have high emission quantum yields. Ferrocene derivatives are often used as electrochemically active labels due to the accessibility of a large variety of derivatives, and their stability and easy redox tuning. [31,32]

Our key substrates were azidonucleosides, which have been used mostly as intermediates to aminonucleosides, but they also exhibit cytotoxic and antiviral properties [33] and are useful photoaffinity probes. [34] Azidonucleotides are available by several methods. [33] For preparation of 2'-azido-2'-deoxyuridine **3** we used a very convenient method starting from 2,2'-*O*-anhydrouridine. [35] 5'-Azido-5'-deoxythymidine **4** and 5'-azido-5'-deoxyuridine **5** were synthesised from 5'-tosylated intermediates, [36] whilst 3'-azido-3'-deoxythymidine **6** was prepared *via* a Mitsunobu type reaction. [37]

Staudinger reaction of azidonucleosides **3-6** with triphenylphosphine led to iminophosphorane intermediates **7-10**, which reacted subsequently with active esters of coumarin-4-acetic acids **2a-c** (Scheme 1) to form amide bond of new nucleoside derivatives **11-14** (Figure 2). [see Supporting Information File 1] It is known that the Staudinger ligation is accelerated in polar solvents, [21] thus we made our experiments in a mixture of acetonitrile and dioxane. The overall reaction rate was high enough at temperatures around 0°C in all cases (1-2 hours reaction time), but at temperatures around -20°C reactions were too slow. The conversion of coumarin-4-acetic acid active esters was almost quantitative in all cases, according to TLC.

The yield of desired products in Staudinger ligations depends on the structure of the carboxylic acid and on the azido group position on the nucleoside. Addition of water to the proposed intermediates **I-IV** (Scheme 2) resulted not only in the formation of the desired products, but also in the formation of








Figure 2: Coumarin labeled nucleosides prepared by intermolecular Staudinger ligation.



Scheme 1: Preparation of coumarin labeled nucleosides, (i) PPh₃, acetonitrile, (ii) HOBT, DCC, dioxane

4-methyl coumarins **15a-c** and aminonucleosides depending on the substrates used. The aminonucleosides and derivatives **15a-c**, as products of a concurrent reaction, complicated the monitoring and the work-up of the reaction mixture. The R_f values of the aminonucleosides and iminophosphorane derivatives **7-10** are very similar and therefore the progress of reaction was monitored and the conversion of substrates was calculated based on coumarin derivatives **2a-c**.

The relationship between the yield of desired product and the azidonucleoside structure is not straightforward. Starting from 2'-azido-2'-deoxyuridine **3** or 5'-azido-5'-deoxythymidine **4** we prepared desired modified nucleosides **11a-c** and **12a-c** in good yields (Table 1). 4-Methyl derivatives of the corresponding coumarin-4-acetic acids were only minor byproducts in these reactions. The isolated yields of polar products were negatively influenced by lengthy separation by flash chromatography. The structures of the products were confirmed by ¹H and ¹³C NMR, ¹H-¹H COSY and ¹³C-¹H HSQC analysis. The signal due to the NH-function of the newly created amide bond was clearly visible in all cases. We expected analogous results in the reactions of 5'-azidonucleoside with coumarin derivatives **2a-c** gave products in low yields, max. 15%. We isolated derivatives **15a-c** as the main products in these cases. According to RP HPLC analyses the reaction mixture contained also 5'-amino-5'-deoxyuridine along with small amount of starting compounds **1a-c**.



Scheme 2: Hydrolysis of proposed intermediates **I(a-c)-IV(a-c)**

The structure of products **13a-c** was confirmed by ¹H NMR analysis. The ratio of product and 4-methyl derivative is probably influenced by diverse effects of nucleobase on the reaction mechanism, especially the interaction of nucleobases with triphenylphosphine residue in the intermediates **I(a-c)-IV(a-c)**. Similar behaviour was observed in the case of 3'-azido-3'-deoxythymidine **6**. Reaction led to the exclusive formation of 4-methyl derivatives **15a-c** and 3'-amino-3'-deoxythymidine,

Table 1: Isolated yields and spectral characteristics of coumarin labeled nucleosides **11–12**^a

Compound ^b	Yield (%)	$\lambda_{ex}(nm)$	$\epsilon (cm^{-1}M^{-1})$	$\lambda_{em}(nm)$	FI (A.U.)
1a	-	326	12487	392	1955
11a	64	326	9981	395	2202
12a	48	325	7123	392	1923
1b	-	323	3032	418	2260
11b ^c	59	324	9033	428	2420
12b	41	322	10006	429	2373
1c	-	318/349	7444	417	719/930
11c	56	319/350	4713	418	822/1006
12c	44	319/350	5291	416	876/1224

^a products **13a–c** were isolated in low yields (max. 15%) and products **14a–c** were not isolated


^b c = 10⁻⁴ M in methanol

^c c = 0.5 × 10⁻⁴ M in methanol

and changes to the standard reaction conditions did not induce the formation of the desired 3'-labeled nucleosides **14a–c**.

The spectral characteristics of the isolated products are summarised in Table 1. We found that all newly prepared conjugates display only one peak in the fluorescence spectrum in methanol. The process of conjugation did not cause a shift in the absorption and fluorescence maxima of our compounds, in comparison with maxima of **1a–c**. We observed only slight changes of fluorescence intensity in the series of coumarin labeled nucleosides.

We also tested the Staudinger ligation as a prospective method for the electrochemical labeling of nucleosides and nucleotides with ferrocene derivatives. 2'-Azido-2'-deoxyuridine **3** was chosen as an azidosubstrate for its satisfactory reactivity in previous experiments and the 2' position of carbohydrate residues is a versatile site for chemical modification of nucleosides. The ferrocene derivative (4-ferrocenyl-4-oxobutanoic acid) **17** was used as an electrochemical label. Analogous reaction of 4-ferrocenyl-4-oxobutanoic acid reactive ester with iminophosphorane nucleoside derivative **7** gave 2'-ferrocene labeled uridine (Scheme 3). The yield of pure isolated product **18** was 57% and its structure was confirmed by ¹H and ¹³C NMR, ¹H-¹H COSY and ¹³C-¹H HSQC analysis.

**Scheme 3:** Preparation of ferrocene labeled uridine

Conclusion

We successfully applied the Staudinger ligation to the synthesis of labeled nucleosides. For this purpose, we prepared several types of azidonucleoside substrates and carboxylic acid derivatives. Although the conversion of reactants was almost quantitative in all cases, the yield of Staudinger ligation was found to be dependent on the structure of the azidonucleoside and the carboxylic acids. A detailed study of the kinetics of the Staudinger ligation with nucleoside substrates, and the possible application to the construction of labeled oligomers and novel bioconjugates for enzyme assays is in progress. A very useful way to use Staudinger ligation in parallel syntheses could be the application of solid-phase reagents to simplify lengthy purification of products.

Supporting Information

Supporting Information File 1

Experimental Section

[<http://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-2-23-S1.pdf>]

Acknowledgments

The present work has been supported by the Slovak Grant Agency VEGA 1/3559/06, Science and Technology Assistance Agency under the contracts No. APVV-51-046505, partly No. APVT-20-031904 and Comenius University Grant No. UK/198/2006. We thank Assoc. Prof. Marta Salisova for the generous gift of the ferrocene substrate and for helpful discussions.

References

1. Chu, C. K., Ed. *Antiviral Nucleosides: Chiral Synthesis and Chemotherapy*, 1st ed.; Elsevier Science: Amsterdam, 2003.

2. De Clercq, E. *Curr. Opin. Microbiol.* **2005**, *8*, 552–560. doi:10.1016/j.mib.2005.08.010
3. De Clercq, E. *Antiviral Res.* **2005**, *67*, 56–75. doi:10.1016/j.antiviral.2005.05.001
4. Waggoner, A. *Curr. Opin. Chem. Biol.* **2006**, *10*, 62–66. doi:10.1016/j.cbpa.2006.01.005
5. Wells, M. *Curr. Opin. Biotechnol.* **2006**, *17*, 28–33. doi:10.1016/j.copbio.2005.12.008
6. Chudakov, D. M.; Lukyanov, S.; Lukyanov, K. A. *Trends Biotechnol.* **2005**, *23*, 605–613. doi:10.1016/j.tibtech.2005.10.005
7. Demidov, V. V. *Trends Biotechnol.* **2003**, *21*, 4–7. doi:10.1016/S0167-7799(02)00008-2
8. Gwynne, P. *Drug, Discovery Development* **2002**, *5*, 50–55.
9. Wojczewski, C.; Stolze, K.; Engels, J. W. *Synlett* **1999**, 1667–1678.
10. Spassova, M.; Kois, P.; Watanabe, K. *Collect. Czech. Chem. Commun.* **1996**, *61*, S290–S293.
11. Wlasoff, W. A.; King, G. C. *Nucleic Acids Res.* **2002**, *30*, e58. doi:10.1093/nar/gnf058
12. Rosi, N. L.; Mirkin, C. A. *Chem. Rev.* **2005**, *105*, 1547–1562. doi:10.1021/cr030067f
13. Brase, S.; Gil, C.; Knepper, K.; Zimmermann, V. *Angew. Chem., Int. Ed.* **2005**, *44*, 5188–5240. doi:10.1002/anie.200400657
14. Staudinger, H.; Meyer, J. *Helv. Chim. Acta* **1919**, *2*, 635–646. doi:10.1002/hlca.19190020164
15. Gololobov, Y. G.; Zhmurova, I. N.; Kasukhin, L. F. *Tetrahedron* **1981**, *37*, 437–472. doi:10.1016/S0040-4020(01)92417-2
16. Gololobov, Y. G.; Kasukhin, L. F. *Tetrahedron* **1992**, *48*, 1353–1406. doi:10.1016/S0040-4020(01)92229-X
17. Saxon, E.; Bertozzi, C. R. *Science* **2000**, *287*, 2007–2010. doi:10.1126/science.287.5460.2007
18. Saxon, E.; Luchansky, S. J.; Hang, H. C.; Yu, C.; Lee, S. C.; Bertozzi, C. R. *J. Am. Chem. Soc.* **2002**, *124*, 14893–14902. doi:10.1021/ja027748x
19. Lemieux, G. A.; de Graffenried, C. L.; Bertozzi, C. R. *J. Am. Chem. Soc.* **2003**, *125*, 4708–4709. doi:10.1021/ja029013y
20. Kohn, M.; Breinbauer, R. *Angew. Chem., Int. Ed.* **2004**, *43*, 3106–3116. doi:10.1002/anie.200401744
21. Lin, F. L.; Hoyt, H. M.; Van Halbeek, H.; Bergman, R. G.; Bertozzi, C. R. *J. Am. Chem. Soc.* **2005**, *127*, 2686–2695. doi:10.1021/ja044461m
22. Soellner, M. B.; Dickson, K. A.; Nilsson, B. L.; Raines, R. T. *J. Am. Chem. Soc.* **2003**, *125*, 11790–11791. doi:10.1021/ja036712h
23. Kohn, M.; Wacker, R.; Peters, C.; Schroder, H.; Souler, L.; Breinbauer, R.; Niemeyer, C. M.; Waldmann, H. *Angew. Chem., Int. Ed.* **2003**, *42*, 5830–5834. doi:10.1002/anie.200352877
24. Wang, C. C.-Y.; Seo, T. S.; Li, Z.; Ruparel, H.; Ju, J. *Bioconjugate Chem.* **2003**, *14*, 697–701. doi:10.1021/bc0256392
25. Nilsson, B. L.; Hondal, R. J.; Soellner, M. B.; Raines, R. T. *J. Am. Chem. Soc.* **2003**, *125*, 5268–5269. doi:10.1021/ja029752e
26. Kiick, K. L.; Saxon, E.; Tirrell, D. A.; Bertozzi, C. R. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, *99*, 19–24. doi:10.1073/pnas.012583299
27. Kovacs, L.; Osz, E.; Domokos, V.; Holzer, W.; Gyorgydeak, Z. *Tetrahedron* **2001**, *57*, 4609–4621. doi:10.1016/S0040-4020(01)00380-5
28. The Handbook, A Guide to Fluorescent Probes and Labeling Technologies. Invitrogen Corporation: Carlsbad, CA, USA, 2006; <http://www.probes.invitrogen.com/handbook> (accessed Mar 10, 2008).
29. deSilva, A. P.; Gunaratne, H. Q. N.; Gunnlaugsson, T.; Huxley, A. J. M.; McCoy, C. P.; Rademacher, J. T.; Rice, T. E. *Chem. Rev.* **1997**, *97*, 1515–1566. doi:10.1021/cr960386p
30. Trenor, S. R.; Shultz, A. R.; Love, B. J.; Long, T. E. *Chem. Rev.* **2004**, *104*, 3059–3077. doi:10.1021/cr030037c
Reference cited therein.
31. van Staveren, D. R.; Metzler-Nolte, N. *Chem. Rev.* **2004**, *104*, 5931–5985. doi:10.1021/cr0101510
32. Zatsepin, T. S.; Andreev, S. Y.; Oretskaya, T. S.; Hianik, T. *Russ. Chem. Rev.* **2003**, *72*, 537–554. doi:10.1070/RC2003v072n06ABEH000807
33. Pathak, T. *Chem. Rev.* **2002**, *102*, 1623–1667. doi:10.1021/cr0104532
References cited therein.
34. Fleming, S. A. *Tetrahedron* **1995**, *51*, 12479–12520. doi:10.1016/0040-4020(95)00598-3
References cited therein.
35. Kirschenheuter, G. P.; Zhai, Y.; Pieken, W. A. *Tetrahedron Lett.* **1994**, *35*, 8517–8520. doi:10.1016/S0040-4039(00)78425-5
36. Ciuffreda, P.; Loseto, A.; Santaniello, E. *Tetrahedron* **2002**, *58*, 5767–5771. doi:10.1016/S0040-4020(02)00575-6
37. Czerniecki, S.; Valery, J.-M. *Synthesis* **1991**, 239–240. doi:10.1055/s-1991-26434

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The license is subject to the Beilstein Journal of Organic Chemistry terms and conditions: (<http://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at:
doi:10.1186/1860-5397-2-23